



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Reports of States and yearly and monthly reports of cities.

ALABAMA—*Mobile*.—Month of June, 1897. Estimated population, 37,817. Total deaths, 77, including phthisis pulmonalis, 7, and enteric fever, 4.

CALIFORNIA—*San Francisco*.—Month of July, 1897. Estimated population, 360,000. Total deaths, 470, including phthisis pulmonalis, 70; enteric fever, 3; diphtheria, 2; croup, 3, and whooping cough, 2.

ILLINOIS—*Chicago*.—Month of July, 1897. Estimated population, 1,750,000. Total deaths, 2,275, including phthisis pulmonalis, 184; enteric fever, 27; scarlet fever, 5; diphtheria, 44; measles, 12, and whooping cough, 21.

MARYLAND—*Baltimore*.—Month of July, 1897. Estimated population—white, 431,054; colored, 75,344; total, 506,398. Deaths, white, 725; colored, 236; total, 961, including phthisis pulmonalis, 68; enteric fever, 13; scarlet fever, 4; diphtheria, 8; measles, 1, and whooping cough, 5.

MASSACHUSETTS—*Brockton*.—Month of July, 1897. Estimated population, 35,853. Total deaths, 38, including phthisis pulmonalis, 4; diphtheria and croup, 1, and whooping cough, 1.

Worcester.—Month of July, 1897. Estimated population, 103,086. Total deaths, 172, including phthisis pulmonalis, 22; enteric fever, 1; scarlet fever, 1, and diphtheria, 3.

MICHIGAN.—Week ended August 14, 1897. Reports to the State board of health, Lansing, from 62 observers, indicate that enteric fever increased and intermittent fever and inflammation of kidney decreased in area of prevalence. Phthisis pulmonalis was reported present during the week at 191, measles at 33, enteric fever at 29, diphtheria at 16, scarlet fever at 14, and whooping cough at 10 places.

Grand Rapids.—Month of July, 1897. Estimated population, 90,000. Total deaths, 71, including phthisis pulmonalis, 5; enteric fever, 1, and diphtheria, 2.

MISSOURI—*Kansas City*.—Month of July, 1897. Estimated population, 175,000. Total deaths, 187, including phthisis pulmonalis, 8; enteric fever, 1; croup, 1, and whooping cough, 1.

NEW YORK—*Yonkers*.—Month of July, 1897. Estimated population, 40,000. Total deaths, 80, including phthisis pulmonalis, 4, and diphtheria, 4.

OHIO—*Cleveland*.—Month of July, 1897. Estimated population, 350,000. Total deaths, 610, including phthisis pulmonalis, 15; enteric fever, 7; scarlet fever, 1; diphtheria, 5, and measles, 8.